arXiv:1605.01294v2 [math.NT] 21 May 2016

ON THE REDUCIBILITY OF SOME SPECIAL QUADRINOMIALS

YONG ZHANG AND HUILIN ZHU

ABSTRACT. Let $n > m > k$ be positive integers and let a, b, c be nonzero rational numbers. We consider the reducibility of some special quadrinomials $x^n + ax^m + bx^k + c$ with $n = 4$ and 5, which related to the study of rational points on certain elliptic curves or hyperelliptic curves.

1. INTRODUCTION

Let $f(x)$ be trinomial or quadrinomial with $deg(f) \geq 3$ and rational coefficients, there are many authors investigated the factorizations of $f(x)$, i.e.,

$$f(x) = f_1(x)f_2(x)\cdots f_k(x), \tag{1.1}$$

where $f_i(x) \in \mathbb{Q}[x]$ with $1 \leq deg(f_i) < deg(f)$, $i = 1, \cdots, k$. We can refer to [2, 3, 5, 6, 7, 8, 9, 10, 11, 12].

A polynomial $f(x)$ with rational coefficients is *primitive reducible* if it is reducible but $f(x^{1/l})$ is not reducible for any integer $l \geq 2$. At the West Coast Number Theory conference in 2007, P.G. Walsh [13] posed "Is there a primitive reducible polynomial of the form $x^i + x^j + x^k + 4$ with $0 < k < j < i$ and $i > 17$?" and "Is there a primitive reducible polynomial of the form $x^i + x^j + x^k + a$ with $0 < k < j < i$, $a \in \mathbb{Z}$ and $a > 4$, and not divisible by a linear or quadratic polynomial?"

Let $f(a, x) = x^n + x^m + x^k + a, n > m > k \geq 1, a \in \mathbb{Q}$. In 2010, J. Jankauskas [7] proved the only primitive quadrinomial $f(4, x)$, such that $f(4, x^l)$ is reducible for some $l > 1$, is $x^4 + x^3 + x^2 + 4$. He also obtained some examples of reducible quadrinomial $x^n + x^m + x^k + a$ with $a \in \mathbb{Z}$ and $a < -5$, such that all the irreducible factors of $f(a, x)$ are of degree ≥ 3. In 2015, A. Bremner and M. Ulas [3] studied the reducible of quadrinomials $f(a, x)$ with $4 \leq n \leq 6$ in a more systematic way and gave further examples of reducible $f(a, x), a \in \mathbb{Q}$, such that all the irreducible factors with degree ≥ 3.

Now we investigate the divisibility of the following quadrinomial

$$f_{n,m,k}(a, b, c, x) = x^n + ax^m + bx^k + c, n > m > k \geq 1, n > 3,$$

by the quadratic polynomial $x^2 + px + q, p^2 + q^2 \neq 0$, where a, b, c are nonzero rational numbers and p, q are rational numbers. In fact, M. Fried and A. Schinzel [5] studied the (ir)reducibility of general quadrinomials in 1972. Here, we care about the cases with a, b, c are special forms such that this problem can be reduced

2010 *Mathematics Subject Classification.* Primary 13P05; Secondary 11G30, 11G05.

Key words and phrases. quadrinomial, elliptic curve, hyperelliptic curve, Chabauty's method.

The first author was supported by China National Natural Science Foundation Grant (No. 11501052). The second author was partly supported by China National Natural Science Foundation Grant (No. 11501477), the Fundamental Research Funds for the Central Universities and the Science Fund of Fujian Province (No. 2015J01024).

to the study of rational points on certain elliptic curves or hyperelliptic curves with genus 1 or 2. Noting that the reducibility of $f_{n,m,k}(a,b,c,x)$ is invariant under the transformations $f(x) \mapsto x^n f(1/x)$ and $f(x) \mapsto f(x)/c$, we just consider the cases

$$x^n + ax^m + x^k + 1, x^n + ax^m + ax^k + 1, x^n + ax^m + x^k + a, x^n + x^m + ax^k + a$$

of $f_{n,m,k}(a,b,c,x)$ for $n = 4$ and 5.

2. Reducible quadrinomials with degree 4

In this section we study the reducibility of the quadrinomial $f_{4,m,k}(a,b,c,x) = x^4 + ax^m + bx^k + c$ for four special cases $(a,b,c) = (a,1,1), (a,a,1), (a,1,a), (1,a,a)$. For the briefness of the discussion, we introduce the following two lemmas.

Lemma 2.1. ([3, Lemma 3.1]) *Let $n \in \mathbb{N}$. Then we have*

$$x^n \mod (x^2 + px + q) = A_n(p,q)x + B_n(p,q),$$

where $A_0(p,q) = 0, A_1(p,q) = 1, B_0(p,q) = 1, B_1(p,q) = 0$, and where for $n \geq 2$:

$$A_n(p,q) = -pA_{n-1}(p,q) - qA_{n-2}(p,q),$$
$$B_n(p,q) = -pB_{n-1}(p,q) - qB_{n-2}(p,q).$$

By the Corollary 3.2 of [3], we have

Lemma 2.2. *If $f_{n,m,k}(a,b,c,x)$ is divisible by x^2+px+q for some a,b,c,p,q, then*

$$\begin{aligned} &A_n(p,q) + aA_m(p,q) + bA_k(p,q) = 0, \\ &c = -B_n(p,q) - aB_m(p,q) - bB_k(p,q). \end{aligned} \tag{2.1}$$

By the theory of elliptic curves, we have the following theorems.

Theorem 2.3. *Let $f_{4,m,k}(a,x) = x^4 + ax^m + x^k + 1$, where $4 > m > k \geq 1$ and $a \neq 0$.*

(1) *If $(m,k) = (2,1)$ then $f_{4,2,1}(a,x)$ is divisible by $x^2 + px + q$ if and only if*

$$a = \frac{q^6 - q^4 - q^3 - q^2 + 1}{q(q-1)^2(q+1)^2}, p = \frac{-q}{q^2-1},$$

where $q \neq 0, \pm 1$. In this case we have

$$f_{4,2,1}\left(\frac{q^6 - q^4 - q^3 - q^2 + 1}{q(q-1)^2(q+1)^2}, x\right) = \left(x^2 - \frac{q}{q^2-1}x + q\right)\left(x^2 + \frac{q^2}{q^2-1}x + \frac{1}{q}\right).$$

(2) *If $(m,k) = (3,1)$ then $f_{4,3,1}(a,x)$ is divisible by $x^2 + px + q$ if and only if*

$$a = 1, p = 2, q = 1; \text{ or } a = 1, p = -1, q = 1.$$

In this case we have

$$f_{4,3,1}(1,x) = (x^2 - x + 1)(x+1)^2.$$

(3) *If $(m,k) = (3,2)$ then $f_{4,3,2}(a,x)$ does not have a quadratic factor.*

Proof of Theorem 2.3. (1) Case $(m,k) = (2,1)$. From Eq. (2.1), we have

$$-p^3 - ap + 2pq + 1 = 0, -p^2q - aq + q^2 + 1 = 0.$$

Solve the above two Diophantine equations with respect to a, we get

$$-\frac{p^3 - 2pq - 1}{p} = -\frac{p^2q - q^2 - 1}{q}.$$

Then

$$p = \frac{-q}{q^2-1}.$$

Hence,

$$a = \frac{q^6 - q^4 - q^3 - q^2 + 1}{q(q-1)^2(q+1)^2}.$$

(2) Case $(m,k) = (3,1)$. From Eq. (2.1), we have

$$ap^2 - p^3 - aq + 2pq + 1 = 0, apq - p^2q + q^2 + 1 = 0.$$

Solve the above two Diophantine equations with respect to a, we get

$$\frac{p^3 - 2pq - 1}{p^2 - q} = \frac{p^2q - q^2 - 1}{pq}.$$

Then

$$p = \frac{q}{2} \pm \frac{\sqrt{4q^3 + q^2 + 4q}}{2}.$$

To make p be a rational number, set $r^2 = 4q^3 + q^2 + 4q$. This is an elliptic curve equivalent to $Y^2 = X^3 + X^2 + 16X$, the rank of which is 0 and the torsion points are $(X,Y) = (0,0), (4; \pm 12)$. (For the torsion points, we omit the point $\mathcal{O}$ at infinity here and in the following.) Hence, $(p,q) = (0,0), (-1,2;1)$, which lead to $a = 1$.

(3) Case $(m,k) = (3,2)$. From Eq. (2.1), we have

$$ap^2 - p^3 - aq - p + 2pq = 0, apq - p^2q - q + q^2 + 1 = 0.$$

Solve the above two Diophantine equations with respect to a, we get

$$\frac{(p^2 - 2q + 1)p}{p^2 - q} = \frac{p^2q - q^2 + q + 1}{pq}.$$

Then

$$p = \pm\sqrt{q(q^2 - q + 1)}.$$

Let $r^2 = q(q^2 - q + 1)$, then p will be a rational number. The above curve is an elliptic curve of rank 0 with trivial torsion points $(r,q) = (0,0), (\pm 1; 1)$. Hence, $(p,q) = (0,0), (\pm 1; 1)$, which lead to $a = 0$. □

Theorem 2.4. *Let $f_{4,m,k}(a,a,1,x) = x^4 + ax^m + ax^k + 1$, where $4 > m > k \geq 1$ and $a \neq 0$.*

(1) *If $(m,k) = (2,1)$ then the set, say $\mathcal{A}$, of those a such that $f_{4,2,1}(a,a,1,x)$ is divisible by $x^2 + px + q$ is infinite. More precisely the set $\mathcal{A}$ is parameterized by the rational points on the rank one elliptic curve*

$$E_1 : \ Y^2 = X^3 + 4X^2 + 16X + 64.$$

(2) *If $(m,k) = (3,1)$ then $f_{4,3,1}(a,a,1,x)$ is divisible by $x^2 + px + q$ if and only if*

$$a = \frac{p^2 - 2}{p}, q = 1,$$

where $p \neq 0$. In this case we have

$$f_{4,3,1}\left(\frac{p^2-2}{p}, \frac{p^2-2}{p}, 1, x\right) = (x^2 + px + 1)(x^2 - \frac{2}{p}x + 1).$$

(3) *If $(m,k) = (3,2)$ it is the case (1) by the transformation $f(x) \mapsto x^4 f(1/x)$.*

Proof of Theorem 2.4. (1) Case $(m,k)=(2,1)$. By Eq. (2.1), we have

$$-p^3-ap+2pq+a=0, -p^2q-aq+q^2+1=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$-\frac{p(p^2-2q)}{p-1}=-\frac{p^2q-q^2-1}{q}.$$

This leads to

$$p^2q-pq^2-q^2+p-1=0.$$

Solve it for q, we have

$$q=\frac{p^2+\sqrt{p^4+4p^2-4}}{2(p+1)}.$$

If q is a rational number, it needs $r^2=p^4+4p^2-4$. This curve is equivalent to the elliptic curve $E_1:\ Y^2=X^3+4X^2+16X+64$, the rank of which is 1 and there are infinitely many rational points on it. Hence, there are infinitely many (p,q), leading to infinitely many a.

(2) Case $(m,k)=(3,1)$. By Eq. (2.1), we have

$$ap^2-p^3-aq+2pq+a=0, apq-p^2q+q^2+1=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$\frac{p(p^2-2q)}{p^2-q+1}=\frac{p^2q-q^2-1}{pq}.$$

This leads to

$$(q-1)(p^2+q^2+1)=0.$$

Then $q=1$. Hence,

$$a=\frac{p^2-2}{p}.$$

The result follows. □

Theorem 2.5. *Let $f_{4,m,k}(a,1,a,x)=x^4+ax^m+x^k+a$, where $4>m>k\geq 1$ and $a\neq 0$.*

(1) If $(m,k)=(2,1)$ then the set, say $\mathcal{A}$, of those a such that $f_{4,2,1}(a,1,a,x)$ is divisible by x^2+px+q is infinite. More precisely the set $\mathcal{A}$ is parameterized by the rational points on the rank one elliptic curve

$$E_1:\ Y^2=X^3+4X^2+16X+64.$$

(2) If $(m,k)=(3,1)$ then $f_{4,3,1}(a,1,a,x)$ is divisible by x^2+px+q if and only if

$$a=p-1, q=p-1,$$

where $p\neq 0,1$. In this case we have

$$f_{4,3,1}(p-1,1,p-1,x)=(x^2-x+1)(x^2+px+p-1).$$

(3) If $(m,k)=(3,2)$ it is the case (1) by the transformations $f(x)\mapsto x^4f(1/x)$ and $f(x)\mapsto f(x)/a$.

Proof of Theorem 2.5. (1) Case $(m,k)=(2,1)$. From Eq. (2.1), we have

$$-p^3-ap+2pq+1=0, -p^2q-aq+q^2+a=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$-\frac{p^3-2pq-1}{p}=-\frac{q(p^2-q)}{q-1}.$$

This leads to

$$p^3+pq^2-2pq+q-1=0.$$

Solve it for q, we have

$$q=\frac{2p-1+\sqrt{-4p^4+4p^2+1}}{2p}.$$

If q is a rational number, it needs $r^2=-4p^4+4p^2+1$. Let $u=\frac{r}{p^2}, v=\frac{1}{p}$, then $u^2=v^4+4v^2-4$. The remainder proof is similar to the case (1) of Theorem 2.4.

(2) Case $(m,k)=(3,1)$. From Eq. (2.1), we have

$$ap^2-p^3-aq+2pq+1=0, apq-p^2q+q^2+a=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$\frac{p^3-2pq-1}{p^2-q}=\frac{q(p^2-q)}{pq+1}.$$

This leads to

$$(-q-1+p)(p^2+pq+q^2+p-q+1)=0.$$

Then $q=p-1$. Hence,

$$a=p-1.$$

The result follows. □

Theorem 2.6. *Let $f_{4,m,k}(1,a,a,x)=x^4+x^m+ax^k+a$, where $4>m>k\geq 1$ and $a\neq 0$.*

(1) *If $(m,k)=(2,1)$ then $f_{4,2,1}(1,a,a,x)$ does not have a quadratic factor.*

(2) *If $(m,k)=(3,1)$ then $f_{4,3,1}(1,a,a,x)$ is divisible by x^2+px+q if and only if*

$$a=-p^3, q=p^2;\ \text{or}\ a=(p-1)^3, q=p-1,$$

where $p\neq 0,1$. In this case we have

$$f_{4,3,1}(1,-p^3,-p^3,x)=(x^2+px+p^2)(x^2-(p-1)x-p);$$

or

$$f_{4,3,1}(1,(p-1)^3,(p-1)^3,x)=(x^2+px+p-1)(x^2-(p-1)x+p^2-2p+1).$$

(3) *If $(m,k)=(3,2)$ it is the case (1) by the transformations $f(x)\mapsto x^4f(1/x)$ and $f(x)\mapsto f(x)/a$.*

Proof of Theorem 2.6. (1) Case $(m,k)=(2,1)$. By Eq. (2.1), we have

$$-p^3+2pq+a-p=0, -p^2q+q^2+a-q=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$p^3-2pq+p=p^2q-q^2+q.$$

This leads to

$$p^3 - p^2q - 2pq + q^2 + p - q = 0.$$

Solve it for q, we have

$$q = \frac{p^2 + 2p + 1 + \sqrt{p^4 + 6p^2 + 1}}{2}.$$

If q is a rational number, it needs $r^2 = p^4 + 6p^2 + 1$. This curve is equivalent to the elliptic curve $Y^2 = X^3 + 6X^2 - 4X - 24$, the rank of which is 0 and the torsion points are $(X, Y) = (-6, 0), (\pm 2, 0)$. Hence, $p = 0, q = 0, 1$, and $a = 0$.

(2) Case $(m, k) = (3, 1)$. By Eq. (2.1), we have

$$-p^3 + p^2 + 2pq + a - q = 0, -p^2q + pq + q^2 + a = 0.$$

Solve the above two Diophantine equations with respect to a, we get

$$p^3 - p^2 - 2pq + q = p^2q - pq - q^2.$$

This leads to

$$(-q - 1 + p)(p^2 - q) = 0.$$

Then $q = p^2$ or $p - 1$. Hence,

$$a = -p^3 \text{ or } (p - 1)^3.$$

The result follows. □

3. Reducible quadrinomials with degree 5

Now we investigate the reducibility of the quadrinomial $f_{5,m,k}(a, b, c, x) = x^5 + ax^m + bx^k + c$ for three special cases $(a, b, c) = (a, 1, 1), (a, a, 1), (a, 1, a), (1, a, a)$. In the following theorems, the cases without asterisk are absolute, and the cases with asterisk represent the results are conjectural.

Theorem 3.1. *Let $f_{5,m,k}(a, x) = x^5 + ax^m + x^k + 1$, where $5 > m > k \geq 1$ and $a \neq 0$.*

(1) *If $(m, k) = (2, 1)$ then $f_{5,2,1}(a, x)$ is divisible by $x^2 + px + q$ if and only if*

$$a = -3, p = -2, q = 1.$$

In this case we have

$$f_{5,2,1}(-3, x) = (x - 1)^2(x^3 + 2x^2 + 3x + 1).$$

(2) *If $(m, k) = (3, 1)$ then $f_{5,3,1}(a, x)$ is divisible by $x^2 + px + q$ if and only if*

$$a = 2, p = -1, q = 1.$$

In this case we have

$$f_{5,2,1}(2, x) = (x^2 - x + 1)(x^3 + x^2 + 2x + 1).$$

(3) *If $(m, k) = (3, 2)$ then $f_{5,3,2}(a, x)$ is divisible by $x^2 + px + q$ if and only if*

$$a = 1, p = -1, q = 1; \text{ or } a = 1, p = 0, q = 1;$$
$$\text{or } a = -\frac{19397}{1458}, p = \frac{10}{27}, q = \frac{1}{6}; \text{ or } a = \frac{2597}{192}, p = -\frac{3}{8}, q = \frac{1}{6}.$$

In this case we have

$$f_{5,3,2}(1,x)=(x+1)(x^2+1)(x^2-x+1);$$
$$\text{or } f_{5,3,2}\Big(-\frac{19397}{1458},x\Big)=\Big(x^2+\frac{10}{27}x+\frac{1}{6}\Big)\Big(x^3-\frac{10}{27}x^2-\frac{40}{3}x+6\Big);$$
$$\text{or } f_{5,3,2}\Big(\frac{2597}{192},x\Big)=\Big(x^2-\frac{3}{8}x+\frac{1}{6}\Big)\Big(x^3+\frac{3}{8}x^2+\frac{27}{2}x+6\Big).$$

*(4) *If $(m,k)=(4,1)$ then $f_{5,4,1}(a,x)$ is divisible by x^2+px+q if and only if*

$$a=-2,p=-1,q=1;\ \text{or } a=-\frac{1055}{16},p=\frac{1}{16},q=\frac{1}{8}.$$

In this case we have

$$f_{5,4,1}(-2,x)=(x^2-x-1)(x^3-x^2-1);$$
$$\text{or } f_{5,4,1}\Big(-\frac{1055}{16},x\Big)=\Big(x^2+\frac{1}{16}x+\frac{1}{8}\Big)(x^3-66x^2+4x+8).$$

*(5) *If $(m,k)=(4,2)$ then $f_{5,4,2}(a,x)$ does not have a quadratic factor.*

(6) *If $(m,k)=(4,3)$ then $f_{5,4,3}(a,x)$ is divisible by x^2+px+q if and only if*

$$a=-1,p=0,q=1.$$

In this case we have

$$f_{5,4,1}(-1,x)=(x^2+1)(x^3-x^2+1).$$

Proof of Theorem 3.1. (1) Case $(m,k)=(2,1)$. From Eq. (2.1), we have

$$p^4-3p^2q-ap+q^2+1=0,p^3q-2pq^2-aq+1=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$\frac{p^4-3p^2q+q^2+1}{p}=\frac{p^3q-2pq^2+1}{q}.$$

This leads to

$$-\frac{p^2q^2-q^3+p-q}{pq}=0.$$

Solve it for p, we have

$$p=\frac{-1\pm\sqrt{4q^5+4q^3+1}}{2q^2}.$$

If p is a rational number, it needs $r^2=4q^5+4q^3+1$. This is a hyperelliptic quintic curve of genus 2. The rank of the Jacobian variety is 1, and Magma's Chabauty routines [4] determine the only finite rational points are $(r,q)=(\pm1;0),(\pm3;1)$, which lead to $(p,q)=(-2,1),(1,1),(0,0)$. Hence, $a=-3$.

(2) Case $(m,k)=(3,1)$. From Eq. (2.1), we have

$$p^4+ap^2-3p^2q-aq+q^2+1=0,p^3q+apq-2pq^2+1=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$-\frac{p^4-3p^2q+q^2+1}{p^2-q}=-\frac{p^3q-2pq^2+1}{pq}.$$

Then

$$\frac{pq^3+p^2-pq-q}{(p^2-q)pq}=0.$$

Solve it for p, we have

$$p=\frac{-q^3+q\pm\sqrt{q^6-2q^4+q^2+4q}}{2}.$$

Let $r^2=q^6-2q^4+q^2+4q$, then p is a rational number. This is a hyperelliptic sextic curve of genus 2. The rank of the Jacobian variety is 1, and Magma's Chabauty routines determine the only finite rational points are $(r,q)=(0,0),(\pm 2;1)$, which lead to $(p,q)=(\pm 1;1),(0,0)$. Hence, $a=2$.

(3) Case $(m,k)=(3,2)$. From Eq. (2.1), we have

$$p^4+ap^2-3p^2q-aq-p+q^2=0, p^3q+apq-2pq^2-q+1=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$-\frac{p^4-3p^2q+q^2-p}{p^2-q}=-\frac{p^3q-2pq^2-q+1}{pq}.$$

This leads to

$$\frac{pq^3+p^2+q^2-q}{(p^2-q)pq}=0.$$

Solve it for p, we have

$$p=\frac{-q^3\pm\sqrt{q^6-4q^2+4q}}{2}.$$

If p is a rational number, it needs $r^2=q^6-4q^2+4q$. This is a hyperelliptic sextic curve of genus 2. The rank of the Jacobian variety is 1, and Magma's Chabauty routines determine the only finite rational points are

$$(r,q)=(0,0),(\pm 1;1),\left(\pm\frac{161}{6};\frac{1}{6}\right),$$

which lead to

$$(p,q)=(0,1),(-1,1),\left(\frac{10}{27},\frac{1}{6}\right),\left(-\frac{3}{8},\frac{1}{6}\right).$$

Hence,

$$a=1,-\frac{19397}{1458},\frac{2597}{192}.$$

*(4) Case $(m,k)=(4,1)$. From Eq. (2.1), we have

$$-ap^3+p^4+2apq-3p^2q+q^2+1=0, -ap^2q+p^3q+aq^2-2pq^2+1=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$\frac{p^4-3p^2q+q^2+1}{p(p^2-2q)}=\frac{p^3q-2pq^2+1}{q(p^2-q)}.$$

Then

$$-\frac{q^4+p^3-p^2q-2pq+q^2}{p(p^2-2q)q(p^2-q)}=0.$$

It needs to consider $q^4+p^3-p^2q-2pq+q^2=0$. Let $p=tq$, we get

$$q^2(qt^3-qt^2+q^2-2t+1)=0.$$

Solve it for q, we get

$$q=\frac{-t^3+t^2\pm\sqrt{t^6-2t^5+t^4+8t-4}}{2}.$$

Let $r^2 = t^6 - 2t^5 + t^4 + 8t - 4$, then q is a rational number. This is a hyperelliptic sextic curve of genus 2. The rank of the Jacobian variety is 2, so standard Chabauty arguments do not apply and we are unable to determine explicitly all the rational points. But the obviously rational points are

$$(r, q) = (\pm 2; 1), \left(\pm \frac{1}{8}; \frac{1}{2} \right),$$

which lead to

$$(p, q) = (0, 0), (-1, -1), (1, 1), \left(\frac{1}{16}, \frac{1}{8} \right).$$

Hence,

$$a = -2, -\frac{1055}{16}.$$

*(5) Case $(m, k) = (4, 2)$. From Eq. (2.1), we have

$$-ap^3 + p^4 + 2apq - 3p^2q - p + q^2 = 0, -ap^2q + p^3q + aq^2 - 2pq^2 - q + 1 = 0.$$

Solve the above two Diophantine equations with respect to a, we get

$$\frac{p^4 - 3p^2q + q^2 - p}{p(p^2 - 2q)} = \frac{p^3q - 2pq^2 - q + 1}{q(p^2 - q)}.$$

This leads to

$$-\frac{q^4 + p^3 + pq^2 - 2pq}{p(p^2 - 2q)q(p^2 - q)} = 0.$$

It needs to consider $q^4 + p^3 + pq^2 - 2pq = 0$. Let $p = tq$, we get

$$q^2(qt^3 + q^2 + qt - 2t) = 0.$$

Solve it for q, we get

$$q = \frac{-t^3 - t \pm \sqrt{t^6 + 2t^4 + t^2 + 8t}}{2}.$$

If q is a rational number, it needs $r^2 = t^6 + 2t^4 + t^2 + 8t$. This is a hyperelliptic sextic curve of genus 2. The rank of the Jacobian variety is 1, but we cannot find the generator of its Jacobian, and Magma's Chabauty routines don't determine the complete rational point. But it is believed that the only rational point is $(r, t) = (0, 0)$, which leads to $(p, q) = (0, 0)$. Hence, the result follows.

(6) Case $(m, k) = (4, 3)$. From Eq. (2.1), we have

$$-ap^3 + p^4 + 2apq + p^2 - 3p^2q - q + q^2 = 0, -ap^2q + p^3q + aq^2 + pq - 2pq^2 + 1 = 0.$$

Solve the above two Diophantine equations with respect to a, we get

$$\frac{p^4 - 3p^2q + p^2 + q^2 - q}{p(p^2 - 2q)} = \frac{p^3q - 2pq^2 + pq + 1}{q(p^2 - q)}.$$

Then

$$-\frac{q^4 + p^3 - q^3 - 2pq}{p(p^2 - 2q)q(p^2 - q)} = 0.$$

It needs to consider $q^4 + p^3 - q^3 - 2pq = 0$. Let $p = tq$, we get

$$q^2(qt^3 + q^2 - q - 2t) = 0.$$

Solve it for q, we get

$$q = \frac{-t^3 + 1 \pm \sqrt{t^6 - 2t^3 + 8t + 1}}{2}.$$

Let $r^2 = t^6 - 2t^3 + 8t + 1$, then q is a rational number. This is a hyperelliptic sextic curve of genus 2. The rank of the Jacobian variety is 1, and Magma's Chabauty routines determine the only finite rational points are $(r, t) = (\pm 1; 1)$, which lead to $(p, q) = (0; 0, 1)$. Hence, $a = -1$. □

Theorem 3.2. *Let $f_{5,m,k}(a, a, 1, x) = x^5 + ax^m + ax^k + 1$, where $5 > m > k \geq 1$ and $a \neq 0$.*

(1) *If $(m, k) = (2, 1)$ then $f_{5,2,1}(a, a, 1, x)$ is divisible by $x^2 + px + q$ if and only if*

$$a = \frac{q^4 + q^3 + q^2 + q + 1}{q}, p = q + 1,$$

where $q \neq 0$. In this case we have

$$f_{5,2,1}\Big(\frac{q^4 + q^3 + q^2 + q + 1}{q}, \frac{q^4 + q^3 + q^2 + q + 1}{q}, 1, x\Big) = (x^2 + (q + 1)x + q)\big(x^3 + (-q - 1)x^2 + (q^2 + q + 1)x + \frac{1}{q}\big).$$

*(2) *If $(m, k) = (3, 1)$ then $f_{5,3,1}(a, a, 1, x)$ does not have a quadratic factor.*

(3) *If $(m, k) = (3, 2)$ then $f_{5,3,2}(a, a, 1, x)$ is divisible by $x^2 + px + q$ if and only if*

$$a = -\frac{q^4 + q^3 + q^2 + q + 1}{q^2}, p = q + 1;$$

or

$$a = \frac{q^4 + q^3 + q^2 + q + 1}{q(q^2 + 2q + 1)}, p = -\frac{q}{q + 1},$$

where $q \neq 0, -1$. In this case we have

$$f_{5,3,2}\Big(-\frac{q^4 + q^3 + q^2 + q + 1}{q^2}, -\frac{q^4 + q^3 + q^2 + q + 1}{q^2}, 1, x\Big) = (x^2 + (q + 1)x + q)\Big(x^3 - (q + 1)x^2 - \frac{(1 + q)x}{q^2} + \frac{1}{q}\Big);$$

or

$$f_{5,3,2}\Big(\frac{q^4 + q^3 + q^2 + q + 1}{q(q^2 + 2q + 1)}, \frac{q^4 + q^3 + q^2 + q + 1}{q(q^2 + 2q + 1)}, 1, x\Big) = \Big(x^2 - \frac{qx}{q + 1} + q\Big)\Big(x^3 + \frac{qx^2}{q + 1} + \frac{x}{q(q + 1)} + \frac{1}{q}\Big).$$

(4) *If $(m, k) = (4, 1)$ then $f_{5,4,1}(a, a, 1, x)$ is divisible by $x^2 + px + q$ if and only if*

$$a = \frac{q^4 + q^3 + q^2 + q + 1}{q(q^2 + q + 1)}, p = q + 1.$$

In this case we have

$$f_{5,4,1}\Big(\frac{q^4 + q^3 + q^2 + q + 1}{q(q^2 + q + 1)}, \frac{q^4 + q^3 + q^2 + q + 1}{q(q^2 + q + 1)}, 1, x\Big) = (x^2 + (q + 1)x + q)\Big(x^3 + \frac{-q^3 - q^2 + 1}{q(q^2 + q + 1)}x^2 + \frac{(q^3 - q - 1)}{q(q^2 + q + 1)}x + \frac{1}{q}\Big).$$

(5) *If* $(m,k)=(4,2)$ *it is the case (2) by the transformations* $f(x)\mapsto x^5f(1/x)$ *and* $f(x)\mapsto f(x)/a$.

(6) *If* $(m,k)=(4,3)$ *it is the case (1) by the transformations* $f(x)\mapsto x^5f(1/x)$ *and* $f(x)\mapsto f(x)/a$.

Proof of Theorem 3.2. (1) Case $(m,k)=(2,1)$. By Eq. (2.1), we have

$$p^4-3p^2q-ap+q^2+a=0, p^3q-2pq^2-aq+1=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$\frac{p^4-3p^2q+q^2}{p-1}=\frac{p^3q-2pq^2+1}{q}.$$

This leads to

$$(-q-1+p)(p^2q+pq-q^2+q-1)=0.$$

Solve it for p, we have

$$p=q+1;\ or\ \frac{-q\pm\sqrt{4q^3-3q^2+4q}}{2q}.$$

If p is a rational number, it needs $r^2=4q^3-3q^2+4q$. Let $u=4r, v=4q$, we have $u^2=v^3-3v^2+16v$. This is an elliptic curve with rank 0 and torsion point $(v,u)=(0,0)$. Then $q=0$. So we get $p=q+1$ and

$$a=\frac{q^4+q^3+q^2+q+1}{q}.$$

*(2) Case $(m,k)=(3,1)$. By Eq. (2.1), we have

$$p^4+ap^2-3p^2q-aq+q^2+a=0, p^3q+apq-2pq^2+1=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$-\frac{p^4-3p^2q+q^2}{p^2-q+1}=-\frac{p^3q-2pq^2+1}{pq}.$$

Then

$$p^3q+pq^3-2pq^2+p^2-q+1=0.$$

This is an equation about p,q with degree 4 and genus 3, so it's hard to get the all rational solutions of it. But we conjecture its rational solution is $(p,q)=(0,1)$.

(3) Case $(m,k)=(3,2)$. By Eq. (2.1), we have

$$p^4+ap^2-3p^2q-ap-aq+q^2=0, p^3q+apq-2pq^2-aq+1=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$-\frac{p^4-3p^2q+q^2}{p^2-p-q}=-\frac{p^3q-2pq^2+1}{q(p-1)}.$$

This leads to

$$(q-1)(pq+p+q)(-q+p-1)=0.$$

Solve it for p, we get

$$p=q+1;\ or\ -\frac{q}{q+1}.$$

Then

$$a=-\frac{q^4+q^3+q^2+q+1}{q^2};\ or\ \frac{q^4+q^3+q^2+q+1}{q(q^2+2q+1)}.$$

(4) Case $(m,k)=(4,1)$. By Eq. (2.1), we have

$$-ap^3+p^4+2apq-3p^2q+q^2+a=0, -ap^2q+p^3q+aq^2-2pq^2+1=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$\frac{p^4-3p^2q+q^2}{p^3-2pq-1}=\frac{p^3q-2pq^2+1}{q(p^2-q)}.$$

Then

$$(q-1)(-q-1+p)(p^2+pq+q^2+p+1)=0.$$

Solve it for p, we get

$$p=q+1;\ or\ \frac{q+1\pm\sqrt{-3q^2+2q-3}}{2}.$$

Noting that $-3q^2+2q-3<0$ for any rational number q, then $p=q+1$. So

$$a=\frac{q^4+q^3+q^2+q+1}{q(q^2+q+1)}.$$

The result follows. □

Theorem 3.3. *Let $f_{5,m,k}(a,1,a,x)=x^5+ax^m+x^k+a$, where $5>m>k\geq 1$ and $a\neq 0$.*

(1) *If $(m,k)=(2,1)$ then $f_{5,2,1}(a,1,a,x)$ does not have a quadratic factor.*

*(2) *If $(m,k)=(3,1)$ then $f_{5,3,1}(a,1,a,x)$ is divisible by x^2+px+q if and only if*

$$a=\frac{1}{2}, p=q=1.$$

In this case we have

$$f_{5,3,2}\big(\frac{1}{2},1,\frac{1}{2},x\big)=(x^2+x+1)\big(x^3-x^2+\frac{x}{2}+\frac{1}{2}\big).$$

(3) *If $(m,k)=(3,2)$ then $f_{5,3,2}(a,1,a,x)$ is divisible by x^2+px+q if and only if*

$$a=-q^2, p=q+1,$$

where $q\neq 0$. In this case we have

$$f_{5,3,2}(-q^2,1,-q^2,x)=(x^2+(q+1)x+q)(x^3-(q+1)x^2+(q+1)x-q).$$

(4) *If $(m,k)=(4,1)$ then $f_{5,4,1}(a,1,a,x)$ does not have a quadratic factor.*

(5) *If $(m,k)=(4,2)$ it is the case (2) by the transformations $f(x)\mapsto x^5f(1/x)$ and $f(x)\mapsto f(x)/a$.*

(6) *If $(m,k)=(4,3)$ it is the case (1) by the transformations $f(x)\mapsto x^5f(1/x)$ and $f(x)\mapsto f(x)/a$.*

Proof of Theorem 3.3. (1) Case $(m,k)=(2,1)$. From Eq. (2.1), we have

$$p^4-3p^2q-ap+q^2+1=0, p^3q-2pq^2-aq+a=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$\frac{p^4-3p^2q+q^2+1}{p}=\frac{pq(p^2-2q)}{q-1}.$$

This leads to

$$p^4+p^2q^2-3p^2q-q^3+q^2-q+1=0.$$

Solve it for p, we have

$$p = \pm\frac{\sqrt{-2q^2+6q+2\sqrt{q^4-2q^3+5q^2+4q-4}}}{2}.$$

If p is a rational number, it needs $r^2 = q^4-2q^3+5q^2+4q-4$ and $s^2 = -2q^2+6q+2r$. Let us consider $r^2 = q^4-2q^3+5q^2+4q-4$, this equation is equivalent to the elliptic curve $Y^2 - 2XY + 8Y = X^3 + 4X^2 + 16X + 64$ with rank 0 and torsion points $(X, Y) = (4, 16), (-4, 0)$. Then $q = \pm 1, p = \pm\sqrt{2}i$. So the result follows.

*(2) Case $(m, k) = (3, 1)$. From Eq. (2.1), we have

$$p^4 + ap^2 - 3p^2q - aq + q^2 + 1 = 0, p^3q + apq - 2pq^2 + a = 0.$$

Solve the above two Diophantine equations with respect to a, we get

$$-\frac{p^4 - 3p^2q + q^2 + 1}{p^2 - q} = -\frac{pq(p^2 - 2q)}{pq + 1}.$$

Then

$$p^4 - pq^3 - 3p^2q + pq + q^2 + 1 = 0.$$

This is an equation about p, q with degree 4 and genus 3, so it's hard to get the all rational solutions of it. But we conjecture its rational solution is $(p, q) = (\pm 1, 1)$. Then $a = \frac{1}{2}$ with $p = q = 1$.

(3) Case $(m, k) = (3, 2)$. From Eq. (2.1), we have

$$p^4 + ap^2 - 3p^2q - aq + q^2 - p = 0, p^3q + apq - 2pq^2 + a - q = 0.$$

Solve the above two Diophantine equations with respect to a, we get

$$-\frac{p^4 - 3p^2q + q^2 - p}{p^2 - q} = -\frac{q(p^3 - 2pq - 1)}{pq + 1}.$$

This leads to

$$p(-q + p - 1)(p^2 + pq + q^2 + p - q + 1) = 0.$$

Solve it for p, we get $p = q + 1$. Then $a = -q^2$.

(4) Case $(m, k) = (4, 1)$. From Eq. (2.1), we have

$$-ap^3 + p^4 + 2apq - 3p^2q + q^2 + 1 = 0, -ap^2q + p^3q + aq^2 - 2pq^2 + a = 0.$$

Solve the above two Diophantine equations with respect to a, we get

$$\frac{p^4 - 3p^2q + q^2 + 1}{p(p^2 - 2q)} = \frac{pq(p^2 - 2q)}{p^2q - q^2 - 1}.$$

Then

$$p^4 + q^4 - 4p^2q + 2q^2 + 1 = 0.$$

Write it as

$$(p^2 - 2q)^2 = -(q^2 - 1)^2,$$

it is easy to see that there is no rational solution (p, q). □

Theorem 3.4. *Let $f_{5,m,k}(1, a, a, x) = x^5 + x^m + ax^k + a$, where $5 > m > k \geq 1$ and $a \neq 0$.*

(1) If $(m, k) = (2, 1)$ then $f_{5,2,1}(1, a, a, x)$ is divisible by $x^2 + px + q$ if and only if

$$a = -(p^2 - p + 1)(p - 1)^2, q = p - 1;$$

or

$$a=\frac{p(p+1)(p^2+p+1)^2}{(2p+1)^2}, q=\frac{p(p^2+p+1)}{2p+1}.$$

In this case we have

$$\begin{aligned}&f_{5,3,2}(1,-(p^2-p+1)(p-1)^2,-(p^2-p+1)(p-1)^2,x)\\&\quad=(x^2+px+p-1)(x^3-px^2-(-p^2+p-1)x-p^3+2p^2-2p+1);\end{aligned}$$

or

$$\begin{aligned}&f_{5,3,2}\bigg(1,\frac{p(p+1)(p^2+p+1)^2}{(2p+1)^2},\frac{p(p+1)(p^2+p+1)^2}{(2p+1)^2},x\bigg)\\&\quad=\bigg(x^2+px+\frac{p(p^2+p+1)}{2p+1}\bigg)\bigg(x^3-px^2+\frac{p^3-p}{2p+1}x+\frac{p^3+2p^2+2p+1}{2p+1}\bigg).\end{aligned}$$

(2) *If* $(m,k)=(3,1)$ *then* $f_{5,3,1}(1,a,a,x)$ *is divisible by* x^2+px+q *if and only if*

$$a=\frac{125}{12}, p=-\frac{1}{2}, q=\frac{5}{18}.$$

In this case we have

$$f_{5,3,2}\bigg(1,\frac{125}{12},\frac{125}{12},x\bigg)=\bigg(x^2-\frac{x}{2}+\frac{5}{18}\bigg)\bigg(x^3+\frac{x^2}{2}+\frac{35x}{36}+\frac{25}{72}\bigg).$$

(3) *If* $(m,k)=(3,2)$ *then* $f_{5,3,2}(1,a,a,x)$ *is divisible by* x^2+px+q *if and only if*

$$a=-p^2, q=p^2,$$

where $p\neq 0$. *In this case we have*

$$f_{5,3,2}(1,-p^2,-p^2,x)=(x-p)(x^2+1)(x^2+px+p^2).$$

(4) *If* $(m,k)=(4,1)$ *then* $f_{5,4,1}(1,a,a,x)$ *is divisible by* x^2+px+q *if and only if*

$$a=\frac{p^4}{2}, q=\frac{p^2}{2};\ \textit{or}\ a=-(p-1)^4, q=p-1.$$

In this case we have

$$f_{5,4,1}\bigg(1,\frac{p^4}{2},\frac{p^4}{2},x\bigg)=(x+1)\bigg(x^2-px+\frac{p^2}{2}\bigg)\bigg(x^2+px+\frac{p^2}{2}\bigg);$$

or

$$f_{5,4,1}(1,-(p-1)^4,-(p-1)^4,x)=(x^2+px+p-1)(x-p+1)(x^2+p^2-2p+1).$$

(5) *If* $(m,k)=(4,2)$ *it is the case (2) by the transformations* $f(x)\mapsto x^5f(1/x)$ *and* $f(x)\mapsto f(x)/a$.

(6) *If* $(m,k)=(4,3)$ *it is the case (1) by the transformations* $f(x)\mapsto x^5f(1/x)$ *and* $f(x)\mapsto f(x)/a$.

Proof of Theorem 3.4. (1) Case $(m,k)=(2,1)$. By Eq. (2.1), we have

$$p^4-3p^2q+q^2+a-p=0, p^3q-2pq^2+a-q=0.$$

Solve the above two Diophantine equations with respect to a, we get

$$-p^4+3p^2q-q^2+p=-p^3q+2pq^2+q.$$

This leads to

$$-(-q-1+p)(p^3+p^2-2pq+p-q)=0.$$

Solve it for q, we have

$$q = p + 1;\ or\ \frac{p(p^2+p+1)}{2p+1}.$$

Then

$$a = -(p^2-p+1)(p-1)^2;\ or\ \frac{p(p+1)(p^2+p+1)^2}{(2p+1)^2}.$$

(2) Case $(m,k) = (3,1)$. By Eq. (2.1), we have

$$p^4 - 3p^2q + p^2 + q^2 + a - q = 0, p^3q - 2pq^2 + pq + a = 0.$$

Solve the above two Diophantine equations with respect to a, we get

$$-p^4 + 3p^2q - p^2 - q^2 + q = -p^3q + 2pq^2 - pq.$$

Then

$$-p^4 + p^3q + 3p^2q - 2pq^2 - p^2 + pq - q^2 + q = 0.$$

Solve it for q, we get

$$q = \frac{p^3+3p^2+p+1+\sqrt{p^6-2p^5+7p^4+3p^2+2p+1}}{2p+1}.$$

If q is a rational number, it needs $r^2 = p^6 - 2p^5 + 7p^4 + 3p^2 + 2p + 1$. This is a hyperelliptic sextic curve of genus 2. The rank of the Jacobian variety is 1, and Magma's Chabauty routines determine the only finite rational points are

$$(r,p) = (\pm 1; 0), \Big(\pm\frac{9}{8}; -\frac{1}{2}\Big),$$

which lead to

$$(p,q) = \big(-\frac{1}{2}, \frac{5}{18}\big).$$

Hence,

$$a = \frac{125}{12}.$$

(3) Case $(m,k) = (3,2)$. By Eq. (2.1), we have

$$p^4 - 3p^2q - ap + p^2 + q^2 - q = 0, p^3q - 2pq^2 - aq + pq + a = 0.$$

Solve the above two Diophantine equations with respect to a, we get

$$\frac{p^4 - 3p^2q + p^2 + q^2 - q}{p} = \frac{pq(p^2 - 2q + 1)}{q-1}.$$

This leads to

$$(p^2 + q^2 - 2q + 1)(p^2 - q) = 0.$$

Solve it for q, we get $q = p^2$. Then $a = -p^3$.

(4) Case $(m,k) = (4,1)$. By Eq. (2.1), we have

$$p^4 - p^3 - 3p^2q + 2pq + q^2 + a = 0, p^3q - p^2q - 2pq^2 + q^2 + a = 0.$$

Solve the above two Diophantine equations with respect to a, we get

$$-p^4 + p^3 + 3p^2q - 2pq - q^2 = -p^3q + p^2q + 2pq^2 - q^2.$$

Then

$$-p(-q - 1 + p)(p^2 - 2q) = 0.$$

Solve it for q, we get

$$q = \frac{p^2}{2};\ or\ p - 1.$$

Then

$$a = \frac{p^4}{2};\ or\ -(p-1)^4.$$

The result follows. □

4. Some related questions

In the end, we raise some questions about the reducibility of quadrinomials.

Question 4.1. Are there infinitely many $f(x) = x^n + ax^m + bx^k + c$, with $abc \neq 0, n > m > k > 0, n \geq 6$ has a quadratic factor $g(x) = x^2 + px + q$ for $(a, b, c) = (a, 1, 1), (a, a, 1), (a, 1, a), (1, a, a)$?

Question 4.2. If $f(x) = x^n + ax^m + bx^k + c, abc \neq 0, n > m > k > 0$, whether all the factors of $f(x)$ have the forms $f_i(x) = x^{n_i} + a_i x^{m_i} + b_i x^{k_i} + c_i$ where $n_i > m_i > k_i > 0, a_i b_i c_i \neq 0$?

Question 4.3. Can we use the method of infinite descent to prove the only rational point on the hyperelliptic curve $r^2 = p^6 + 2p^4 + p^2 + 8p$ is $(0, 0)$?

References

1. W. Bosma, J. Cannon and C. Playoust, The Magma algebra system. I. The user language, J. Symbolic Comput. **24** (1997), 235–265.
2. A. Bremner and M. Ulas, On the reducibility type of trinomials, Acta Arith. **153** (2012), 349–372.
3. A. Bremner and M. Ulas, Some observations concerning reducibility of quadrinomials, Acta Math. Hungar. **145** (2015), 320–349.
4. Chabauty's Method, http://magma.maths.usyd.edu.au/magma/handbook/text/1468.
5. M. Fried and A. Schinzel, Reducibility of quadrinomials, Acta Arith. **21** (1972), 153–171.
6. L. Hajdu and R. Tijdeman, Polynomials dividing infinitely many quadrinomials or quintinomials, Acta Arith. **107** (2003), 381–404.
7. J. Jankauskas, On the reducibility of certain quadrinomials, Glas. Mat. Ser. III **45 (65)** (2010), 31–41.
8. W. Ljunggren, On the irreducibility of certain trinomials and quadrinomials, Math. Scand. **8** (1960), 65–70.
9. A. Schinzel, On Reducible Trinomials, Dissertationes Mathematicae **329** (1993), 83 pp.
10. A. Schinzel, On reducible trinomials, II, Publ. Math. Debrecen **56** (2000), 575–608.
11. A. Schinzel, On reducible trinomials, III, Period. Math. Hung. **43** (2001), 43–69.
12. A. Schinzel, Selecta, Volume I: Diophantine problems and polynomials. Edited by Henryk Iwaniec, Władysław Narkiewicz and Jerzy Urbanowicz. Heritage of European Mathematics. Zürich: European Mathematical Society Publishing House (EMS), 2007.
13. West Coast Number Theory Conference, Problem Set, http://westcoastnumbertheory.org/problem-sets/.

School of Mathematics and Statistics, Changsha University of Science and Technology, Changsha 410114, People's Republic of China

E-mail address: zhangyongzju@163.com

School of Mathematical Sciences, Xiamen University, Xiamen 361005, People's Republic of China

E-mail address: hlzhu@xmu.edu.cn